

UC San Diego

UC San Diego Previously Published Works

Title

The debut of chiral cyclic (alkyl)(amino)carbenes (CAACs) in enantioselective catalysis.

Permalink

<https://escholarship.org/uc/item/5tr3t9q2>

Journal

Chemical science, 10(33)

ISSN

2041-6520

Authors

Pichon, Delphine
Soleilhavoup, Michele
Morvan, Jennifer
et al.

Publication Date

2019-09-01

DOI

10.1039/c9sc02810b

Peer reviewed



The debut of chiral cyclic (alkyl)(amino)carbenes (CAACs) in enantioselective catalysis†

Delphine Pichon, ^a Michele Soleilhavoup, ^b Jennifer Morvan, ^a
Glen P. Junor, ^b Thomas Vives, ^a Christophe Crévisy, ^a Vincent Lavallo, ^b
Jean-Marc Campagne, ^c Marc Mauduit, ^{*a} Rodolphe Jazzar ^{*b}
and Guy Bertrand ^{*b}

The popularity of NHCs in transition metal catalysis has prompted the development of chiral versions as electron-rich neutral stereodirecting ancillary ligands for enantioselective transformations. Herein we demonstrate that cyclic (alkyl)(amino)carbene (CAAC) ligands can also engage in asymmetric transformations, thereby expanding the toolbox of available chiral carbenes.


DOI: 10.1039/c9sc02810b

rsc.li/chemical-science

the first examples of chiral CAAC ligands in asymmetric catalysis.


In our initial paper on CAACs,⁹ we showed that the enantiopure *L*-Menth⁺CAAC could be prepared without time-consuming enantio- or diastereoselective separation from the inexpensive (–)-menthol (Scheme 2). The key step of the synthesis was based on the well-known propensity of relatively bulky reactants to approach the cyclohexane moiety selectively from the equatorial direction.

Years ago, we tested L -MenthCAAC transition metal complexes in a variety of asymmetric catalytic reactions without any success. Given our recent mechanistic work on copper-catalysis, we decided to revive this topic. We considered benchmarking the L -MenthCAAC in the copper-catalysed Asymmetric Conjugate Borylation (ACB) reaction. Over the past decade, this chemical transformation has emerged as a stalwart method for the preparation of chiral organoboron building blocks, which are



Scheme 1 Synthetic route to cyclic (alkyl)(amino)carbenes, and existing families of CAACs.



† Electronic supplementary information (ESI) available: Experimental procedures, NMR spectra. CCDC 1919315. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c9sc02810b

Scheme 2 Synthesis of *L*-MenthCAAC highlighting the key step.

sought after in organic synthesis.¹⁸ Although pioneered by Yun¹⁹ and others using chiral phosphine ligands,^{18d} Fernández,²⁰ Hoveyda²¹ and others^{18d} have successfully shown that carbene ligands, specifically NHCs, could also be utilized in this process.

As can be seen in Scheme 3, although the *L*-MenthCAAC copper complex **1** was found to be highly active in the addition of bis(pinacolato)diboron (B_2Pin_2) to various α,β -unsaturated esters, an almost complete lack of asymmetric induction was observed. In comparison, the known non C_2 -chiral NHC copper complex **2**,²² derived from (*S*)-1-(naphthalen-1-yl)ethan-1-amine, gave a 55% ee for the borylated adduct **P₁**. Examining the X-ray crystal structure of **1**,⁹ we expected some asymmetric induction since for many years we believed that a conformational inversion of the menthyl ring was energetically inaccessible. However, we recently found that in the solid state the menthyl group of the *L*-MenthCAAC amine adduct *L*-MenthCAAC^{NH} existed as the other conformer with the methyl and isopropyl substituents in axial position (Scheme 4).²³ We were able to confirm with DFT calculations that the inverted menthyl conformer is also readily accessible in complex **1** (Fig. 1). The existence of the two conformers **1** and **1'** readily explains the lack of asymmetric induction since they have antagonistic stereo-inducing effects.

To circumvent this issue, we sought to prepare a more rigid chiral CAAC. Among the number of readily available molecules derived from the chiral pool, we selected the steroid backbone for its bulk, structural diversity, and tunability. We also reasoned that owing to a three-dimensional skeleton composed of four fused rings (three six-member cyclohexane rings – A, B, C- and one five-member cyclopentane ring D), this steroid motif

Scheme 4 Conformational inversion of the menthyl ring.²³Fig. 1 Proposed justification for the lack of asymmetric induction in catalyst **1**.

would provide the desired rigid structural features while keeping the chirality intact during the synthesis (Fig. 2).

Before proceeding further, we confirmed that our synthetic methodology¹⁶ extends to the readily available decahydronaphthalene, a model substrate mimicking the first two fused cyclohexane (A and B) rings of a steroid skeleton (Scheme 5). CAAC-decahydronaphthyl iminium *rac*-NaphCAAC was obtained in 7 steps and 50% yield as a white powder.

Reassured by these results, we applied the same synthetic strategy to 5 α -cholestan-3-one, an inexpensive enantiopure







Fig. 2 Steroid CAACs as versatile chiral ligand frameworks.



Scheme 3 Comparing *L*-MenthCAACCuCl **1** and NHC-CuCl **2** in the Asymmetric Conjugate Borylation (ACB) reaction. ^aDetermined by ¹H NMR spectroscopy using mesitylene as an internal standard. ^bIsolated yield after SiO₂ purification. ^cDetermined by GC on a chiral stationary phase (see ESI for details†).

Scheme 5 Preparation of *rac*-NaphCAAC.



Scheme 6 Preparation of CholestCAAC and complex 3.

steroid derived from coprostanol (Scheme 6). The CAAC-cholestanyl iminium CholestCAAC was obtained in 6 steps and 63% yield as a white fluffy powder.²⁴ The corresponding copper complex 3 was prepared by deprotonation of the CholestCAAC iminium with KHMDS followed by addition of copper chloride in THF. Single crystals were obtained by slow diffusion of diethyl ether in acetonitrile, and an X-ray diffraction study confirmed the formation of CholestCAACuCl complex 3 (Fig. 3). It crystallized in the $P2_1$ space group with one molecule in the asymmetric unit cell.

The absolute stereochemistry was conclusively established using anomalous dispersion with a Flack parameter of 0.002(7)





Fig. 3 X-ray crystal structure of 3 (top) with the side-on orientation (omitting hydrogen for clarity) and an overlaid topological steric map with a quadrant diagram.



Scheme 7 Evaluation of CholestCAACuCl complex 3 in ACB reaction.

^aDetermined by ¹H NMR spectroscopy using mesitylene as an internal standard. ^bIsolated yield after SiO₂ purification. ^cDetermined by GC on a chiral stationary phase (see ESI for details†). ^dAnthracene was used as internal standard.

from refinement. The X-ray data of 3 show a distinctive orientation of the cholestanyl backbone, which gives rise to the topographic steric map showed in Fig. 3.^{25,26} As can be seen, the corresponding quadrant diagram could support facial stereo-selectivity in catalyst-substrate adducts.

We next evaluated complex 3 in the ACB reaction (Scheme 7). As with *L*-MenthCAACuCl 1, the steroid copper complex CholestCAACuCl 3 efficiently catalyzes the addition of B₂pin₂ to various β-substituted α,β-unsaturated esters, affording the corresponding 1,4-adducts in moderate to good isolated yields (47 to 77%). More importantly, we were pleased to observe enantio-inductions with enantioselectivities reaching 55%.

3. Conclusions

This work rationalizes the inability of *L*-MenthCAAC to induce enantiomeric excess, and indicates that upon providing the right steric environment, the use of chiral CAACs should not be overlooked in metal-catalyzed asymmetric transformations. This new family of stereo-directing chiral carbene ligands are readily available from inexpensive precursors belonging to the chiral pool, and it is noteworthy that the chirality is not restricted to chiral amines (as is the case of NHCs), but extends to chiral aldehydes, a much larger feedstock.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements


This work was supported by the U.S. Department of Energy, Office of Science, Basic Energy Sciences, Catalysis Science Program, under Award # DE-SC0009376, and the Agence Nationale de la Recherche (ANR-16-CE07-0019 – Hel-NHC)



Thanks are due to the Alfred P. Sloan Foundation's University Centre for Exemplary Mentoring and the National Science Foundation Graduate Research Fellowship Program under Grant No. DGE-1650112 (G. P. J.), and to the National Science Foundation (V. L. – Grant # CHE-1455348). Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation. We also greatly acknowledge the Region Bretagne for the ARED 2018, ("Bio-meta", N° 601), and Rennes-Metropole for supporting international PhD mobility (D. P.). MM and TV thank Shimadzu and Chiral Technology for their support in the separation of chiral molecules by Supercritical Fluid Chromatography (SFC) technology.

Notes and references

- (a) J. C. Sheehan and D. H. Hunneman, *J. Am. Chem. Soc.*, 1966, **88**, 3666–3667; (b) D. Enders, H. Gielen, G. Raabe, J. Runsink and J. H. Teles, *Chem. Ber.*, 1996, **129**, 1483–1488; (c) W. A. Herrmann, L. J. Goossen, C. Köcher and G. R. J. Artus, *Angew. Chem., Int. Ed.*, 1996, **35**, 2805–2807; (d) M. T. Powell, D. R. Hou, M. C. Perry, X. Cui and K. Burgess, *J. Am. Chem. Soc.*, 2001, **123**, 8878–8879; (e) T. J. Seiders, D. W. Ward and R. H. Grubbs, *Org. Lett.*, 2001, **3**, 3225–3228.
- For reviews: (a) F. Wang, L.-J. Liu, W. Wang, S. Li and M. Shi, *Coord. Chem. Rev.*, 2012, **256**, 804–853; (b) M. Zhao, Y.-T. Zhang, J. Chen and L. Zhou, *Asian J. Org. Chem.*, 2018, **7**, 54–69; (c) J. Ouyang and J. Crassous, *Coord. Chem. Rev.*, 2018, **376**, 533–547; (d) D. M. Flanagan, F. Romanov-Michailidis, N. A. White and T. Rovis, *Chem. Rev.*, 2015, **115**, 9307–9387; (e) A. Grossmann and D. Enders, *Angew. Chem., Int. Ed.*, 2012, **51**, 314–325.
- Selected examples of chiral NHC in: – Au catalysis – (a) Y.-M. Wang, C. N. Kuzniewski, V. Rauniyar, C. Hoong and F. D. Toste, *J. Am. Chem. Soc.*, 2011, **133**, 12972–12975; (b) Y. Matsumoto, K. B. Selim, H. Nakanishi, K. Yamada, Y. Yamamoto and K. Tomioka, *Tetrahedron Lett.*, 2010, **51**, 404–406; (c) C. Bartolome, D. García-Cuadrado, Z. Ramiro and P. Espinet, *Inorg. Chem.*, 2010, **49**, 9758–9764; (d) S. Handa and L. M. Slaughter, *Angew. Chem., Int. Ed.*, 2012, **51**, 2912–2915; – Cu catalysis – ; (e) S. Drissi-Amraoui, M. Morin, C. Crévisy, O. Baslé, R. Marcia de Figueiredo, M. Mauduit and J.-M. Campagne, *Angew. Chem., Int. Ed.*, 2015, **54**, 11830–11834; – Pd catalysis – ; (f) E. P. Kündig, T. M. Seidel, Y. X. Jia and G. Bernardinelli, *Angew. Chem., Int. Ed.*, 2007, **46**, 8484–8487; – Rh catalysis – ; (g) J. H. Kim, S. Greßies, M. Bouladakis-Arapinis, C. Daniliuc and F. Glorius, *ACS Catal.*, 2016, **6**, 7652–7656. – Ru catalysis – ; (h) A. Kannenberg, D. Rost, S. Eibauer, S. Tiede and S. Blechert, *Angew. Chem., Int. Ed.*, 2011, **50**, 3299–3302.
- D. Janssen-Mueller, C. Schlepphorst and F. Glorius, *Chem. Soc. Rev.*, 2017, **46**, 4845–4854.
- For reviews, see: (a) A. Vivancos, C. Segarra and M. Albrecht, *Chem. Rev.*, 2018, **118**, 9493–9586; (b) G. Guisado-Barrios, M. Soleilhavoup and G. Bertrand, *Acc. Chem. Res.*, 2018, **51**, 3236–3244; (c) M. Melaimi, M. Soleilhavoup and G. Bertrand, *Angew. Chem., Int. Ed.*, 2010, **49**, 8810–8849.
- A. J. Arduengo III, R. L. Harlow and M. Kline, *J. Am. Chem. Soc.*, 1991, **113**, 361–363.
- A. J. Arduengo III, J. R. Goerlich and W. J. Marshall, *J. Am. Chem. Soc.*, 1995, **117**, 11027–11028.
- D. Enders, K. Breuer, G. Raabe, J. Runsink, J. H. Teles, J. P. Melder, K. Ebel and S. Brode, *Angew. Chem., Int. Ed.*, 1995, **34**, 1021–1023.
- V. Lavallo, Y. Canac, C. Präsang, B. Donnadiou and G. Bertrand, *Angew. Chem., Int. Ed.*, 2005, **44**, 5705–5709.
- For reviews on CAACs see: (a) M. Melaimi, R. Jazzar, M. Soleilhavoup and G. Bertrand, *Angew. Chem., Int. Ed.*, 2017, **56**, 10046–10068; (b) M. Soleilhavoup and G. Bertrand, *Acc. Chem. Res.*, 2015, **48**, 256–266; (c) M. Melaimi, M. Soleilhavoup and G. Bertrand, *Angew. Chem., Int. Ed.*, 2010, **49**, 8810–8849; (d) S. Roy, K. C. Mondal and H. W. Roesky, *Acc. Chem. Res.*, 2016, **49**, 357–369; (e) U. S. D. Paul and U. Radius, *Eur. J. Inorg. Chem.*, 2017, 3362–3375.
- (a) V. M. Marx, A. H. Sullivan, M. Melaimi, S. C. Virgil, B. K. Keitz, D. S. Weinberger, G. Bertrand and R. H. Grubbs, *Angew. Chem., Int. Ed.*, 2014, **54**, 1919–1923; (b) D. R. Anderson, V. Lavallo, D. J. O'leary, G. Bertrand and R. H. Grubbs, *Angew. Chem., Int. Ed.*, 2007, **46**, 7262–7265; (c) D. R. Anderson, T. Ung, G. Mkrtumyan, G. Bertrand, R. H. Grubbs and Y. Schrodi, *Organometallics*, 2008, **27**, 563–566; (d) V. M. Marx, A. H. Sullivan, M. Melaimi, S. C. Virgil, B. K. Keitz, D. S. Weinberger, G. Bertrand and R. H. Grubbs, *Angew. Chem., Int. Ed.*, 2015, **54**, 1919–1923.
- C. M. Weinstein, G. P. Junor, D. R. Tolentino, R. Jazzar, M. Melaimi and G. Bertrand, *J. Am. Chem. Soc.*, 2018, **140**, 9255–9260.
- (a) M. P. Wiesenfeldt, Z. Nairoukh, W. Li and F. Glorius, *Science*, 2017, **357**, 908–912; (b) Y. Wei, B. Rao, X. Cong and X. Zeng, *J. Am. Chem. Soc.*, 2015, **137**, 9250–9253; (c) L. Tran, J. L. Fulton, J. C. Linehan, M. Balasubramanian, J. A. Lercher and R. M. Bullock, *ACS Catal.*, 2019, **9**, 4106–4114; (d) Z. Nairoukh, M. Wollenburg, C. Schlepphorst, K. Bergander and F. Glorius, *Nat. Chem.*, 2019, **11**, 264–270.
- (a) E. A. Romero, R. Jazzar and G. Bertrand, *Chem. Sci.*, 2017, **8**, 165–168; (b) J. Chu, D. Munz, R. Jazzar, M. Melaimi and G. Bertrand, *J. Am. Chem. Soc.*, 2016, **138**, 7884–7887.
- (a) X. Hu, D. Martin, M. Melaimi and G. Bertrand, *J. Am. Chem. Soc.*, 2014, **136**, 13594–13597; (b) R. Kinjo, B. Donnadiou and G. Bertrand, *Angew. Chem., Int. Ed.*, 2011, **50**, 5560–5563; (c) X. Zeng, R. Kinjo, B. Donnadiou and G. Bertrand, *Angew. Chem., Int. Ed.*, 2010, **49**, 942–945; (d) V. Lavallo, G. D. Frey, B. Donnadiou, M. Soleilhavoup and G. Bertrand, *Angew. Chem., Int. Ed.*, 2008, **47**, 5224–5228; (e) L. Jin, D. S. Weinberger, M. Melaimi, C. E. Moore, A. L. Rheingold and G. Bertrand, *Angew. Chem., Int. Ed.*, 2014, **53**, 9059–9063.
- (a) R. Jazzar, H. Liang, B. Donnadiou and G. Bertrand, *J. Organomet. Chem.*, 2006, **691**, 3201–3205; (b) R. Jazzar, R. D. Dewhurst, J.-B. Bourg, B. Donnadiou, Y. Canac and



- G. Bertrand, *Angew. Chem., Int. Ed.*, 2007, **46**, 2899–2902; (c) R. Jazzar, J.-B. Bourg, R. D. Dewhurst, B. Donnadiu and G. Bertrand, *J. Org. Chem.*, 2007, **72**, 3492–3499; (d) X. Zeng, G. D. Frey, R. Kinjo, B. Donnadiu and G. Bertrand, *J. Am. Chem. Soc.*, 2009, **131**, 8690–8696.
- 17 E. Tomás-Mendivil, M. M. Hansmann, C. M. Weinstein, R. Jazzar, M. Melaimi and G. Bertrand, *J. Am. Chem. Soc.*, 2017, **139**, 7753–7756.
- 18 (a) J. A. Schiffner, K. Muther and M. Oestreich, *Angew. Chem., Int. Ed.*, 2010, **49**, 1194–1196; (b) E. C. Neeve, S. J. Geier, I. A. Mkhaliid, S. A. Westcott and T. B. Marder, *Chem. Rev.*, 2016, **116**, 9091–9161; (c) Q. Liu, B. Tian, P. Tian, X. Tong and G.-Q. Lin, *Chin. J. Org. Chem.*, 2015, **35**, 1–14; (d) K. Zheng, X. Liu and X. Feng, *Chem. Rev.*, 2018, **118**, 7586–7656.
- 19 (a) J. E. Lee and J. Yun, *Angew. Chem., Int. Ed.*, 2008, **47**, 145–147; (b) H. Chea, H.-S. Sim and J. Yun, *Adv. Synth. Catal.*, 2009, **351**, 855–858; (c) H. S. Sim, X. Feng and J. Yun, *Chem.–Eur. J.*, 2009, **15**, 1939–1943; (d) X. Feng and J. Yun, *Chem. Commun.*, 2009, 6577–6579.
- 20 V. Lillo, A. Prieto, A. Bonet, M. M. Díaz-Requejo, J. Ramírez, P. J. Pérez and E. Fernández, *Organometallics*, 2009, **28**, 659–662.
- 21 J. M. O'Brien, K. S. Lee and A. H. Hoveyda, *J. Am. Chem. Soc.*, 2010, **132**, 10630–10633.
- 22 (a) C. Jahier-Diallo, M. S. T. Morin, P. Queval, M. Rouen, I. Artur, P. Querard, L. Toupet, C. Crévisy, O. Baslé and M. Mauduit, *Chem.–Eur. J.*, 2015, **21**, 993–997. For a higher ee, see: ; (b) J. K. Park, H. H. Lackey, M. D. Rexford, K. Kovnir, M. Shatruk and D. T. McQuade, *Org. Lett.*, 2010, **12**, 5008–5011.
- 23 D. Tolentino, S. Neale, C. Isaac, S. Macgregor, M. Whittlesey, R. Jazzar and G. Bertrand, *J. Am. Chem. Soc.*, 2019, **141**, 9823–9826.
- 24 Cholesterol derived NHCs have been reported: (a) L. Rakers, L. M. Martínez-Prieto, A. M. López-Vinasco, K. Philippot, P. W. N. M. van Leeuwen, B. Chaudret and F. Glorius, *Chem. Commun.*, 2018, **54**, 7070–7073; (b) L. Rakers, D. Grill, A. L. L. Matos, S. Wulff, D. Wang, J. Börgel, M. Körsgen, H. F. Arlinghaus, H.-J. Galla, V. Gerke and F. Glorius, *Cell Chem. Biol.*, 2018, **25**, 952–961.
- 25 H. Clavier and S. P. Nolan, *Chem. Commun.*, 2010, **46**, 841–861.
- 26 Steric maps were extrapolated using the SambVca 2 web-based program developed by Cavallo and co-workers. F. Falivene, R. Credendino, A. Poater, A. Petta, L. Serra, R. Oliva, V. Scarano and L. Cavallo, *Organometallics*, 2016, **35**, 2286–2293.

